



ROMIVEX™ DIP



MITICIDE - INSECTICIDE & REPELLENT

GENERAL INFORMATION

Romivex™ DIP is a contact miticide and insecticide recommended for dip application to control mites and soft-bodied insects, such as two-spotted spider mites, aphids, thrips, mealybugs, and whiteflies on all plants and crops. Romivex DIP has a 0-hour restricted entry interval (REI) and 0-day preharvest interval (PHI).

ACTIVE INGREDIENT

Rosemary oil50%

OTHER INGREDIENTS

Glycerol monooleate, Sorbitol, Lauric acid, Myristic acid, Palmitic acid, Stearic acid, Oleic acid..... 50%

TOTAL:.....100%

**KEEP OUT OF REACH OF CHILDREN
WARNING (AVISO)**



FIRST AID

IF IN EYES: Rinse with water. Get medical attention if irritation develops and persists. **IF ON SKINOR CLOTHING:** Wash off with soap and water. Get medical attention if irritation develops and persists. **IF INHALED:** Move to fresh air. Call a poison control center or doctor for further treatment advice if symptoms develop or persist. **IF INGESTED:** Rinse mouth with clean water. Call a poison control center or doctor if symptoms occur. **HOTLINE NUMBER:** For emergency call the poison control center at 1-800-222-1222. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information concerning this product, call the National Pesticides Information Center at 1-800-858-7378, 6:30 a.m. to 4:30 p.m. Pacific time (PT), seven days a week. For non-emergency Kemin product inquiries, please call Customer Service at 1-800-752-2864 between the hours of 7 a.m. to 4:30 p.m. (Central Standard Time) Monday-Friday.

PRECAUTIONARY STATEMENTS

If you do not understand the label, find someone to explain it to you in detail (Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle). Wear protective gloves, protective clothing, and eye protection or face protection.



Manufactured by:
KEMIN INDUSTRIES, INC.
2100 MAURY ST.
DES MOINES, IOWA U.S.A. 50317

**NET CONTENT:
1 GALLON**



DIRECTIONS FOR USE

READ THE ENTIRE LABEL – USE IN ACCORDANCE WITH LABEL INSTRUCTIONS.

This product is a concentrate and must be diluted before use. Use the dip dilution immediately. Do not store the diluted solution.

USE SITES

For use on all agricultural plants and crops. Indoor and outdoor, not limited to greenhouses, nurseries, ornamentals, trees, shrubs, flowers, landscapes, bedding plants, tropical plants, seedlings, plugs and liners and unrooted cuttings. Romivex™ DIP is not recommended for cut flowers.

RECOMMENDED APPLICATION RATES

A fresh treatment solution should be prepared for every 3,000 cuttings.

0.25 - 0.5% (v/v): For 1 gallon of water: 0.32 fl.oz. - 0.64 fl.oz. For 10 gallon of water: 3.2 fl.oz - 6.4 fl.oz.

DIP APPLICATION INSTRUCTIONS - SHAKE WELL BEFORE USING

Clean and disinfect the dipping containers. Make a new dip treatment solution by diluting the recommended amount of Romivex DIP in sufficient clean water at room temperature to allow complete submersion of the cuttings to be treated. The use of an adjuvant is not required. Submerge the entire cuttings in the dip treatment solution with gentle agitation for about 15 to 60 seconds to wet the front and back of the leaves and provide full coverage of the cuttings. NOTE: The submersion time is sufficient when water stops bubbling. Finally, remove the cuttings from the dip submergence solution and allow the excess to drain as needed. Do not stack the draining cuttings. The cuttings are ready to plant immediately.

ADDITIONAL SPRAY APPLICATION INSTRUCTIONS

Suggested dilution rates 32-256 fl.oz./100 gallon of water. A minimum of 100 gallons of finished solution to treat 40,000 sq.ft. (1 acre) is recommended. Use standard spray equipment. Treatment Interval: Every 5 to 7 days.

COMPATIBILITY OF MIXTURES

Limited compatibility testing has been conducted for Romivex DIP with other commonly used chemistries (i.e. calcium, hormones, fungicides). As such, mixing or use of Romivex DIP with any other products shall be the exclusive risk and responsibility of the user. Consult specific product labels for additional information. Conduct a mix compatibility jar test when you mix Romivex DIP with other products and test the mixture on a small scale before making large-scale applications.

PLANT SAFETY (PHYTOTOXICITY)

Since plant varieties are numerous and may react differently to Romivex DIP, test the product on a small scale before making large-scale applications. Keep cuttings out of direct sunlight and maintain a cool temperature (60-70°F) during treatment. Do not perform dip treatment on plants under stress or when temperatures exceed 90°F.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water or food by storage and disposal. Store in original container in a cool, well-ventilated, dry place (68-80°F). Do not freeze. Keep container closed when not in use. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. **IF EMPTY:** Offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration. **IF PARTLY FILLED:** Call your local waste agency for disposal instructions.

FIFRA 25(b) EXEMPT

This product is a minimum-risk pesticide that has not been registered with the United States Environmental Protection Agency because it qualifies for exemption from registration under Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) Section 25(b).

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.

